

Debreu's social equilibrium theorem with asymmetric information and a continuum of agents

Nicholas C. Yannelis

Received: 21 April 2006 / Revised: 28 March 2007 / Published online: 3 May 2007
© Springer-Verlag 2007

Abstract We provide several different generalizations of Debreu's social equilibrium theorem by allowing for asymmetric information and a continuum of agents. The results not only extend the ones in Kim and Yannelis (J Econ Theory 77:330–353, 1977), Yannelis and Rustichini (Stud Econ Theory 2:23–48, 1991), but also new theorems are obtained which allow for a convexifying effect on aggregation (non-concavity assumption on the utility functions) and non-convex strategy sets (pure strategies). This is achieved by imposing the assumption of “many more agents than strategies” (Rustichini and Yannelis in Stud Econ Theory 1:249–265, 1991; Tourky and Yannelis in J Econ Theory 101:189–221, 2001; Podczeck in Econ Theory 22:699–725, 2003).

Keywords Social equilibrium · Asymmetric information · Many more players than strategies · Convexifying effect · Pure strategy equilibrium

JEL Classification Numbers D82 · D50 · D83 · C62 · C71 · D46 · D61

1 Introduction

Debreu (1952) introduced the idea of a social system (or abstract economy, a generalization of Nash (1950, 1951) normal form game) and proved the existence of an

To the memory of Gerard Debreu. A preliminary draft was presented in Paris, in April of 2005. I have benefited from the discussion, comments and questions of Erik Balder, Jean-Marc Bonnisseu, Bernard Cornet and Filipe Martins Da-Rocha and Conny Podczeck. A careful and knowledgeable referee made several useful comments and rescued me from a mishap.

N. C. Yannelis (✉)
Department of Economics, University of Illinois at Urbana–Champaign,
Champaign, IL 61820, USA
e-mail: nyanneli@uiuc.edu

equilibrium for a social system. This result was the main mathematical tool used to prove the existence of a Walrasian equilibrium for a concrete economy by [Arrow and Debreu \(1954\)](#).

The purpose of this paper is twofold. First, to generalize the idea of a social system by introducing asymmetric information, and a continuum of agents. Second, to prove several existence of an equilibrium theorems for a social system with asymmetric information with a continuum of agents and with an infinite dimensional strategy space.

Our first theorem extends several known results. First, it extends the [Schmeidler \(1973\)](#) theorem to allow for asymmetric information. Second, it extends the [Yannelis \(1991\)](#), [Kim and Yannelis \(1977\)](#) and [Balder \(2002\)](#) existence results from a Bayesian normal form game with a continuum of players to a social Bayesian system with a continuum of agents. Furthermore, it extends the Bayesian social equilibrium existence theorem of [Yannelis \(2002\)](#) from a finite set of agents to a continuum one.

Our second theorem is similar in spirit with our first one, but it does not require the assumption of concavity of the utility function. However, the externalities, i.e., the strategies of all other player affecting the utility of a player, are modeled as an integral rather than as a product, as it is the case in our first theorem. This approach allows for a convexifying effect. For abstract economies or social systems without asymmetric information this type of convexifying effect was introduced by [Haller \(1993\)](#), [see also [Da-Rocha and Topuzu \(2005\)](#)]. It should be noted, that since we work with an infinite dimensional strategy space the Lyapunov theorem fails. Thus, we impose the assumption of “many more agents than strategies” which assures that the Lyapunov theorem still holds (see [Rustichini and Yannelis \(1991\)](#), [Tourky and Yannelis \(2001\)](#) for a detailed analysis of the idea of many more agents than commodities). This assumption is obviously satisfied whenever the strategy space is finite dimensional and there is a continuum of agents.

The first two theorems are proved for a Bayesian decision making framework, i.e., agents maximize interim expected utility and update their priors. We also provide counterparts of the first two theorems for the case of ex ante expected utility maximizing behaviour.

It should be mentioned that recently [Balder \(forthcoming\)](#) and [Cornet and Topuzu \(forthcoming\)](#) have extended the [Aumann \(1965\)](#) Walrasian equilibrium existence theorem to allow for interdependent preferences. Our social equilibrium existence theorems with asymmetric information could be used to extend the recent deterministic results of Balder and Cornet-Topuzu by allowing for asymmetric information.

The mathematics used in this paper is rather diverse. We employ several results on the continuity and measurability of the set of integrable selections from a Banach-valued correspondence as well as results on the semicontinuity of the integral of a Banach-valued correspondence. Furthermore, some compactness and convexity results for the integral of Banach-valued correspondence play an important role.

The paper is organized as follows: The next section contains definitions. Section 3 introduces the model, i.e., the social system with asymmetric information and a continuum of agents. Section 4 contains existence results for interim expected utility functions (Bayesian decision making) and Sect. 5 contains similar results for ex ante expected utility function. Finally, Sect. 6 contains a purification result and some

concluding remarks and open questions are in Sect. 7. Several mathematical results used in the paper are collected in the Appendix for the convenience of the reader.

2 Definitions

Let X and Y be sets. 2^A denotes the set of all nonempty subsets of the set A . \mathcal{R} denotes the set of real numbers. The graph G_ϕ of a correspondence $\phi : X \rightarrow 2^Y$ is the set $G_\phi = \{(x, y) \in X \times Y : y \in \phi(x)\}$. If X and Y are topological spaces, $\phi : X \rightarrow 2^Y$ is said to be *lower-semicontinuous* (l.s.c) if the set $\{x \in X : \phi(x) \cap V \neq \emptyset\}$ is open in X for every open subset V of Y ; $\phi : X \rightarrow 2^Y$ is said to be *upper-semicontinuous* (u.s.c) if the set $\{x \in X : \phi(x) \subset V\}$ is open in X for every open subset V of Y ; $\phi : X \rightarrow 2^Y$ is said to be *continuous* if it is u.s.c. and l.s.c.

If (X, α) and (Y, β) are measurable spaces and $\phi : X \rightarrow 2^Y$ is a correspondence, ϕ is said to have a *measurable graph* if G_ϕ belongs to the product σ -algebra $\alpha \otimes \beta$. We are often interested in the situation where (X, α) is a measurable space, Y is a topological space and $\beta = \beta(Y)$ is the Borel σ -algebra of Y . For a correspondence ϕ from a measurable space into a topological space, if we say that ϕ has a measurable graph, it is understood that the topological space is endowed with its Borel σ -algebra. In the same setting as above, i.e., (X, α) , a measurable space and Y a Banach space, ϕ is said to be *lower measurable* if $\{x \in X : \phi(x) \cap V \neq \emptyset\} \in \alpha$ for every V open in Y . It follows from the projection theorem¹ that if a correspondence has a measurable graph then it is also lower measurable. The reverse is also true if the correspondence is closed valued.

Let (T, τ, μ) be a finite measure space and X be a Banach space. Denote by $L_1(\mu, X)$ the space of equivalence classes of X -valued Bochner integrable functions $x : T \rightarrow X$ normed by

$$\|x\| = \int_T \|x(t)\| d\mu(t).$$

It is a standard result that normed by the functional $\|\cdot\|$ above, $L_1(\mu, X)$ becomes a Banach space [see (Diestel and Uhl 1997, p. 50)]. We denote by S_ϕ^1 the *set of all selections from $\phi : T \rightarrow 2^X$* that belong to the space $L_1(\mu, X)$, i.e.,

$$S_\phi^1 = \{x \in L_1(\mu, X) : x(t) \in \phi(t) \mu - a.e.\}.$$

S_ϕ^1 is the *set of all Bochner integrable selections from $\phi(\cdot)$* . Recall that the correspondence $\phi : T \rightarrow 2^X$ is said to be *integrably bounded* if there exists a map $h \in L_1(\mu, \mathfrak{R})$ such that $\sup\{\|x\| : x \in \phi(t)\} \leq h(t) \mu - a.e$. Moreover, note that if T is a complete measure space, X is a separable Banach space and $\phi : T \rightarrow 2^X$ is an integrably bounded, nonempty valued correspondence having a measurable graph, then by virtue of the Aumann measurable selection theorem (see Appendix) we can conclude that S_ϕ^1 is nonempty. It should be noted that for the applicability of the Aumann

¹ In this case the measure space must be complete.

measurable selection theorem, the range of the correspondence φ , i.e., the space X must be separable and metrizable. For this reason in the subsequent sections we will make sure that $L_1(\mu, X)$ is separable.

The integral of the correspondence $\varphi : T \rightarrow 2^X$ is defined as follows:

$$\int_T \varphi(t) d\mu(t) = \left\{ \int_T x(t) d\mu(t) : x \in S_\varphi^1 \right\}.$$

In the sequel we will denote the above integral by $\int_T \varphi$, or $\int \varphi$.

Observe that, if S_φ^1 is nonempty, we can conclude that $\int_T \varphi$ is also nonempty.

3 The model

3.1 The Debreu social system with asymmetric information

Let $(\Omega, \mathcal{F}, \mu)$ be a complete, finite separable measure space, where Ω , denotes the set of *states of nature of the world* and the σ -algebra \mathcal{F} denotes the set of *events*. Let Y be a separable Banach space, denoting the *commodity* or *strategy* space. A *social system with asymmetric information* and with a measure² *space of agents* (T, τ, ν) , is a set $\Gamma = \{(X, u, A, \mathcal{F}_t, q_t) : t \in T\}$ where,

1. $X : T \times \Omega \rightarrow 2^Y$ is the *random action (strategy)* set-valued function, where, $X(t, \omega) \subset Y$ is interpreted as the strategy set of agent t of the state of nature ω . Define the set $S_{X_t}^1$ as:

$$S_{X_t}^1 = \{y(t) \in L_1(\mu, Y) : y(t, \cdot) : \Omega \rightarrow Y \text{ is } \mathcal{F}_t\text{-measurable and } y(t, \omega) \in X(t, \omega) \mu - a.e.\}.$$

Notice that $S_{X_t}^1$ is the set of all Bochner integrable and \mathcal{F}_t -measurable selections from the random strategy set of agent t . In essence this is the set, out of which agent t will pick his/her optimal choices. In particular, an element x_t in $S_{X_t}^1$ is called a *strategy* for agent t . The typical element of $S_{X_t}^1$ is denoted by \tilde{x}_t and that of $X(t, \omega)$ by $x_t(\omega)$ (or x_t). Let $S_X^1 = \{\tilde{x} \in L_1(\nu, L_1(\mu, Y)) : \tilde{x}(t) \in S_{X_t}^1 \nu - a.e.\}$. An element of S_X^1 will be a *joint strategy profile*.

2. For each fixed $(t, \omega) \in T \times \Omega$, $u(t, \omega, \cdot, \cdot) : L_1(\nu, Y) \times X(t, \omega) \rightarrow \mathcal{R}$ is the *random utility function*, where $u(t, \omega, x, x_t)$ is interpreted as the utility function of agent t , at the state of nature ω , using his/her strategy x_t and all other players use the joint strategy x .
3. $A : T \times \Omega \times S_X^1 \rightarrow 2^Y$, is the *random constraint correspondence* of agent t , where for all $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$, $A(t, \omega, \tilde{x}) \subset X(t, \omega)$, and $A(t, \omega, \tilde{x})$ is interpreted as the constraint of agent t , when the state is ω and other agents use the joint strategy \tilde{x} .

² The measure space (T, τ, ν) is assumed to be complete and finite.

4. \mathcal{F}_t is the sub σ -algebra of \mathcal{F} which denotes the *private information* of agent t .
5. $q_t : \Omega \rightarrow \mathcal{R}_{++}$ is the *prior* of agent t , i.e., a Radon-Nikodym derivative such that $\int_{\Omega} q_t(\omega) d\mu(\omega) = 1$.

3.2 Interim expected utility (Bayesian case)

It will be convenient to assume throughout the paper that Ω is a *countable* set and that the σ -algebra \mathcal{F}_t is generated by a countable partition Λ of Ω . For each $\omega \in \Omega$, let $E_t(\omega)$ in Λ denote the smallest set in \mathcal{F}_t containing ω and to assume that for each t , $\int_{\omega' \in E_t(\omega)} q_t(\omega') d\mu(\omega') > 0$. For each fixed $(t, \omega) \in T \times \Omega$, the *interim expected utility* of agent t , $U(t, \omega, \cdot, \cdot) : S_X^1 \times X(t, \omega) \rightarrow \mathcal{R}$ is defined as

$$U(t, \omega, \tilde{x}, x_t) = \int_{\omega' \in E_t(\omega)} u(t, \omega', \tilde{x}(\omega'), x_t(\omega')) q_t(\omega' | E_t(\omega)) d\mu(\omega')$$

where

$$q_t(\omega' | E_t(\omega)) = \begin{cases} 0 & \text{if } \omega' \notin E_t(\omega) \\ \frac{q_t(\omega')}{\int_{\tilde{\omega} \in E_t(\omega)} q_t(\tilde{\omega}) d\mu(\tilde{\omega})} & \text{if } \omega' \in E_t(\omega). \end{cases}$$

Definition 4.1 A *social equilibrium* for Γ is a strategy profile $\tilde{x} \in S_X^1$ such that for $v - a.e.$

- (1) $\tilde{x}(t, \omega) \in A(t, \omega, \tilde{x}) \mu - a.e.$ and
- (2) $U(t, \omega, \tilde{x}, \tilde{x}(t, \omega)) = \max_{y \in A(t, \omega, \tilde{x})} U(t, \omega, \tilde{x}, y) \mu - a.e.$

A couple of comments are in order: First notice that $\tilde{x} \in S_X^1$ implies that for each fixed $t \in T$, $x(t, \cdot)$ is an \mathcal{F}_t -measurable selection from $X(t, \cdot)$, i.e., $x(t, \cdot)$ is chosen by agent t so that it reflects his/her own private information. Condition (1) indicates that the optimal choice is in the constraint correspondence for almost all agents and almost all states, (i.e., $v - a.e.$ and $\mu - a.e.$). Condition (2) is the best reply notion, i.e., no player can deviate from his/her optimal strategy (in his/her constraint set) and increase his/her payoff once all other agents have chosen the optimal strategy vector \tilde{x} . Again this holds for almost all states and almost all agents, i.e., $\mu - a.e.$ and $v - a.e.$

4 Bayesian social equilibrium existence theorems

4.1 Existence of social asymmetric equilibrium

We begin by stating the assumptions needed to prove our first existence theorem.

Assumptions

(A.1)

- (a) $X : T \times \Omega \rightarrow 2^Y$ is a nonempty, convex, weakly compact valued and integrably bounded, set valued function,

- (a') $X : T \times \Omega \rightarrow 2^Y$ is a nonempty, weakly compact valued and integrably bounded, set valued function.
 - (b) For each fixed $t \in T$, $X(t, \cdot)$ has an \mathcal{F}_t -measurable graph, i.e., $G_{X(t, \cdot)} \in \mathcal{F}_t \otimes \beta(Y)$.
- (A.2)
- (a) For each $(t, \omega) \in T \times \Omega$, $u(t, \omega, \cdot, \cdot) : L_1(v, Y) \times X(t, \omega) \rightarrow \mathcal{R}$ is weakly continuous.
 - (b) For each fixed $(x, y) \in L_1(v, Y) \times Y$, $u(\cdot, \cdot, x, y) : T \times \Omega \rightarrow \mathcal{R}$ is a measurable function.
 - (c) For each $(t, \omega, x) \in T \times \Omega \times L_1(v, Y)$, $u(t, \omega, x, \cdot) : X(t, \omega) \rightarrow \mathcal{R}$ is concave.
 - (d) For each $t \in T$, $u(t, \cdot, \cdot, \cdot)$ is integrably bounded.
- (A.3)
- (a) $A : T \times \Omega \times S_X^1 \rightarrow 2^Y$ has a measurable graph.
 - (b) For each $(t, \omega) \in T \times \Omega$, $A(t, \omega, \cdot) : S_X^1 \rightarrow 2^Y$ is a weakly continuous correspondence with closed, convex and nonempty values.

(A.4)
 The correspondence $t \mapsto S_{X_t}^1$ has a measurable graph.

Theorem 1 *Let Γ be a social system with asymmetric information satisfying (A.1) (a), (b) – (A.4). Then a social equilibrium exists in Γ .*

Proof First observe that the set $S_{X_t}^1$ is nonempty. Indeed in view of assumption A.1(b), by virtue of the Aumann measurable selection theorem, we can conclude that $S_{X_t}^1$ is nonempty, and similarly, we can also conclude that S_X^1 is also nonempty (recall (A.4)). It follows from (A.2) and the fact that Ω is a countable set that for each $(t, \omega) \in T \times \Omega$, $U(t, \omega, \cdot, \cdot) : S_X^1 \times X(t, \omega) \rightarrow \mathcal{R}$ is weakly continuous³. Notice that for each fixed $(t, \tilde{x}, y) \in T \times S_X^1 \times Y$, $U(t, \cdot, \tilde{x}, y) : \Omega \rightarrow \mathcal{R}$ is \mathcal{F}_t -measurable and for each $(\tilde{x}, y) \in S_X^1 \times Y$, $U(\cdot, \cdot, \tilde{x}, y) : T \times \Omega \rightarrow \mathcal{R}$ is $\tau \otimes \mathcal{F}_t$ -measurable. Furthermore, by (A.2) (c) it follows that for $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$, $U(t, \omega, \tilde{x}, \cdot)$ is concave.

Define $F : T \times \Omega \times S_X^1 \rightarrow 2^Y$ by

$$F(t, \omega, \tilde{x}) = \{y \in A(t, \omega, \tilde{x}) : U(t, \omega, \tilde{x}, y) = \max_{z \in A(t, \omega, \tilde{x})} U(t, \omega, \tilde{x}, z)\}.$$

It can be easily checked that F is nonempty valued. Indeed, since all the values of the set valued function A are contained in the weakly compact set $X(\cdot, \cdot)$ and A is closed and convex (hence weakly closed), it follows that A is weakly compact valued. Since, for each $(t, \omega, \tilde{x}) \in T \times \Omega \times S_X^1$, $U(t, \omega, \tilde{x}, \cdot)$ is weakly continuous [see Balder and Yannelis (1993)], we conclude that F is nonempty valued. It follows from (A.2) (c), that F is convex valued.

By virtue of the Berge maximum theorem, for each fixed $(t, \omega) \in T \times \Omega$, $F(t, \omega, \cdot) : S_X^1 \rightarrow 2^Y$ is weakly u.s.c. Furthermore, from (Castaing and Valadier 1997,

³ See Balder and Yannelis (1993) for complete arguments of the continuity of the expected utility.

Lemma III.39, p. 86), $F(t, \cdot, \tilde{x}) : \Omega \rightarrow 2^Y$ has a measurable graph, and $F(\cdot, \cdot, \tilde{x}) : T \times \Omega \rightarrow 2^Y$ has a $\tau \otimes \mathcal{F}$ -measurable graph.

Define the set-valued function $\varphi : T \times S_X^1 \rightarrow 2^{L_1(\mu, Y)}$ by

$$\varphi(t, \tilde{x}) = \{\tilde{y}(t) \in L_1(\mu, Y) : \tilde{y}(t, \omega) \in F(t, \omega, \tilde{x}) \mu - a.e.\} \cap S_{X_t}^1.$$

By the measurability lifting theorem (see Appendix) the correspondence $t \mapsto \{\tilde{y}(t) \in L_1(\mu, Y) : \tilde{y}(t, \omega) \in F(t, \omega, \tilde{x}) \mu - a.e.\}$ has a measurable graph and so does $t \mapsto S_{X_t}^1$ by (A.4). Thus, for each fixed $\tilde{x} \in S_X^1$, $\varphi(\cdot, \tilde{x})$ has a measurable graph. Since for each fixed $\tilde{x} \in S_X^1$, $F(\cdot, \cdot, \tilde{x})$ has a measurable graph and it is nonempty-valued then by the Aumann measurable selection theorem, it admits a measurable selection and we can conclude that φ is nonempty valued. It follows from the convex valueness of F that φ is also convex valued. By Diestel’s theorem (see Appendix) the set $S_{X_t}^1$ is a weakly compact subset of $L_1(\mu, Y)$ and so is S_X^1 . It follows from the u.s.c. lifting theorem (see Appendix) that for each fixed t , $\varphi(t, \cdot)$ is weakly u.s.c.⁴ Define the correspondence $\Phi : S_X^1 \rightarrow 2^{S_X^1}$ by

$$\Phi(\tilde{x}) = \{\tilde{y} \in S_X^1 : \tilde{y}(t) \in \varphi(t, \tilde{x}) \nu - a.e.\}.$$

Another application of the u.s.c. lifting theorem (see Appendix) enables us to conclude that Φ is a weakly u.s.c. correspondence which is obviously convex valued (since φ is convex valued) and also nonempty valued (recall once more the Aumann measurable selection theorem). Since the set S_X^1 is weakly compact, convex and nonempty the Fan-Glicksberg theorem is applicable and there is a fixed point, i.e., there exist $\tilde{x}^* \in S_X^1$ such that $\tilde{x}^* \in \Phi(\tilde{x}^*)$, which implies that for $\nu - a.e.$, (1) $\tilde{x}^*(t, \omega) \in A(t, \omega, \tilde{x}^*) \mu - a.e.$ and (2) $U(t, \omega, \tilde{x}^*, \tilde{x}^*(t, \omega)) = \max_{y \in A(t, \omega, \tilde{x}^*)} U(t, \omega, \tilde{x}^*, y) \mu - a.e.$ This completes the proof of the theorem.

4.2 Convexifying effect

We will now show that the concavity assumption on the utility function can be dropped as well as the convex valueness from the strategy sets. The key condition that it is introduced is the assumption of “many more agents than strategies”, that is, the dimensionality of the set of agents is greater than the dimensionality of the strategy space. In this set up, the set of joint strategies S_X^1 , will now be replaced by the integral of the set-valued strategy correspondence X . As previously, let $S_{X_t}^1 = \{y(t) \in L_1(\mu, Y) : y(t, \cdot) : \Omega \rightarrow Y \text{ is } \mathcal{F}_t - \text{measurable and } y(t, \omega) \in X(t, \omega) \mu - a.e.\}$, and let $S_X^1 = \{\tilde{y} \in L_1(\nu, L_1(\mu, Y)) : \tilde{y}(t) \in S_{X_t}^1 \nu - a.e.\}$. Define the integral of X as: $\int_T X \equiv \int_T X(t) d\nu(t) = \{\int_T \tilde{x} : \tilde{x} \in S_X^1\}$.

The *interim expected utility* of agent t , $\bar{U}(t, \omega, \cdot, \cdot) : \int_T X \times X(t, \omega) \rightarrow \mathcal{R}$ is defined as previously, but now the set of joint strategies has been replaced by the

⁴ I.e., the set $\{x \in S_X^1 : \varphi(t, x) \subset V\}$ is weakly open in S_X^1 for every weakly open subset V of S_X^1 , see also Appendix (u.s.c. lifting theorem).

integral, $\int_T X$. The bar indicates that this interim expected utility has different domain than the one in Sect. 4.1. The idea of the proof is similar. However, we have to replace in the augment, the u.s.c. lifting theorems, with the fact that integration preserves u.s.c., and also make use of the Rustichini and Yannelis (1991) theorem which proves that the integral of a Banach-valued correspondence is convex provided that the dimensionality of the space of agents is bigger than the dimensionality of the strategy space. All these technical details can be found in the Appendix.

Definition 4.2 A (convexifying) social asymmetric equilibrium for Γ is a strategy profile \tilde{x}^* in $\int_T X$, i.e., there exist $\tilde{x}(t) \in S_{X_t}^1 \quad \nu - a.e.$, $\int_T \tilde{x} = \tilde{x}^*$, such that, for $\nu - a.e.$,

- (i) $\tilde{x}(t, \omega) \in A(t, \omega, \tilde{x}^*) \quad \mu - a.e.$, and
- (ii) $\bar{U}(t, \omega, \tilde{x}^*, \tilde{x}(t, \omega)) = \max_{y \in A(t, \omega, \tilde{x}^*)} \bar{U}(t, \omega, \tilde{x}^*, y) \quad \mu - a.e.$

The idea of a convexifying equilibrium is not new, for a different framework it was introduced by Haller (1993).⁵ Haller works with a finite dimensional strategy space, thus, he does have “many more agents than strategies” implicitly in his model. Haller (1993), also considers infinite dimensional strategy spaces but the concavity assumption cannot be dispensed with, because in his framework the Lyapunov theorem fails (i.e., the dimension assumption (A.5), may not hold).

The notion of the equilibrium is the same as in definition 4.1, except that now condition (2) indicates that no agent can deviate from his/her optimal strategy and increase his/her payoff. However, all other player’s strategy is captured by the integral (average sum of the optimal individual strategies), i.e., $\tilde{x}^* = \int_T \tilde{x}$ and the effect of each individual’s strategy $\tilde{x}(t, \cdot)$ on the aggregate (average) is simply zero. This is in essence the idea of perfect competition, which is captured by our model. Notice that in definition 4.1, we are dealing with products and there is no convexifying effect on aggregation.

The assumption of “many more agents than strategies” is formally stated:

- (A.5) The pair $((T, \tau, \nu), K)$, where K is any separable Banach space, satisfies the condition of Theorem A in the Appendix, i.e., if $E \in \tau, \nu(E) > 0$, then $\dim L_{\infty, E}(\nu) > \dim K$.

Theorem 2 Let Γ be a social system with asymmetric information satisfying (A.1), (a’) and (b), (A.2), (a), (b), (d) and (A.3)–(A.5). Then a convexifying social asymmetric equilibrium exists in Γ .

Proof As in the proof of Theorem 1, the continuity and measurability properties of the interim expected utility $\bar{U} : T \times \Omega \times \int_T X \times X(t, \omega) \rightarrow \mathcal{R}$ are still valid, but now for each $(t, \omega, \tilde{x}) \in T \times \Omega \times \int_T X, \bar{U}(t, \omega, \tilde{x}, \cdot)$ is not concave. We will show below that this is not a problem.

Define $F : T \times \Omega \times \int_T X \rightarrow 2^Y$ by

$$F(t, \omega, \tilde{x}) = \{y \in A(t, \omega, \tilde{x}) : \bar{U}(t, \omega, \tilde{x}, y) = \max_{z \in A(t, \omega, \tilde{x})} \bar{U}(t, \omega, \tilde{x}, z)\}.$$

⁵ An early version was circulated as a working paper at VPI in 1986.

As in the Proof of Theorem 1, F is nonempty valued, recall that $\bar{U}(t, \omega, \tilde{x}, \cdot)$ is weakly continuous and A is weakly compact valued so the maximum is obtained. As in the previous theorem, for each fixed $(x, \omega) \in T \times \Omega$, $F(t, \omega, \cdot) : \int_T X \rightarrow 2^Y$ is weakly u.s.c. and for each fixed $\tilde{x} \in \int_T X$, $F(\cdot, \cdot, \tilde{x}) : T \times \Omega \rightarrow 2^Y$ has a measurable graph.

Define the correspondence $\varphi : T \times \int_T X \rightarrow 2^{L_1(\mu, Y)}$ by

$$\varphi(t, \tilde{x}) = \{\tilde{y}(t) \in L_1(\mu, Y) : \tilde{y}(t, \omega) \in F(t, \omega, \tilde{x}) \ \mu - a.e.\}.$$

For each fixed $t \in T$, $\varphi(t, \cdot)$ is weakly u.s.c. and weakly compact valued and for each fixed $\tilde{x} \in \int_T X$, $\varphi(\cdot, \tilde{x})$ has a measurable graph (recall the theorems in the Appendix). Define $\Phi : \int_T X \rightarrow 2^{\int_T X}$ by $\Phi(\tilde{x}) = \int_T \varphi(t, \tilde{x})$. Then Φ is weakly u.s.c., (integration preserves u.s.c. Theorem, see Appendix), nonempty valued (recall Aumann’s measurable selection theorem), and by Theorem A in the Appendix it is convex. As noted in the Appendix, $\int_T X$ is weakly compact and clearly convex, and nonempty. Thus, by the Fan-Glicksberg fixed point theorem there exist $\tilde{x}^* \in \int_T X$ such that $\tilde{x}^* \in \Phi(\tilde{x}^*)$. It can be easily checked that \tilde{x}^* is a convexifying social asymmetric equilibrium for Γ .

5 Ex ante expected utility

Notice that both theorems in Sects. 4.1 and 4.2 are obtained for interim expected utility functions, and this is the Bayesian framework where agents receive a signal as to what is the event in their partition which contains the realized state of nature. Thus, agents condition their expected utility on the event which contains the realized state of nature and update their priors using Bayes’ rule. In this set up the (interim) conditional expected utility depends on the realized state of nature and the arguments required to prove Theorems 1 and 2 are a bit different than the ex ante case we consider below. In the ex ante case there is no signaling and agents integrate their utility function over any possible state of nature. Thus, the (ex ante) expected utility doesn’t depend on the states of nature. In particular, the ex ante expected utility $\bar{\bar{U}} : T \times S_X^1 \times S_{X_t}^1 \rightarrow \mathcal{R}$ is defined as

$$\bar{\bar{U}}(t, \tilde{x}, x_t) = \int_{\Omega} u(t, \tilde{x}(\omega), x_t(\omega))d\mu(\omega).$$

We can now define the corresponding notion of Definition 4.1 for the ex ante expected utility.

Definition 5.3 An (ex ante) social equilibrium for Γ is a strategy profile $\tilde{x} \in S_X^1$ such that for $v - a.e.$,

- (i) $\tilde{x} \in S_A^1(t, \tilde{x}) = \{y \in S_{X_t}^1 : y(t, \omega) \in A(t, \omega, \tilde{x}) \ \mu - a.e.\}$, and
- (ii) $\bar{\bar{U}}(t, \tilde{x}, \tilde{x}_t) = \max_{z \in S_A^1(t, \tilde{x})} \bar{\bar{U}}(t, \tilde{x}, z)$.

Theorem 3 Let Γ be a social system with asymmetric information satisfying (A.1) (a), (b)–(A.4). Then there exists an ex ante social equilibrium in Γ .

Proof The idea of the proof remains essentially the same. Some modifications are required. We outline the argument.

Define $S_A^1 : T \times S_X^1 \rightarrow 2^{L_1(\mu, Y)}$ by

$$S_A^1(t, \tilde{x}) = \{y \in S_{X_t}^1 : y(t, \omega) \in A(t, \omega, \tilde{x}) \ \mu - a.e.\}.$$

By the measurability lifting theorem for each fixed $\tilde{x} \in S_X^1$, $S_A^1(\cdot, \tilde{x})$ has a measurable graph and by Corollary 5.6 in (Yannelis 1991, p. 21), for each fixed $t \in T$, $S_A^1(t, \cdot)$ is (weakly) continuous. It follows from the Aumann measurable selection theorem that S_A^1 is nonempty valued (recall that A has a measurable graph).

Define the correspondence $F : T \times S_X^1 \rightarrow 2^{L_1(\mu, Y)}$ by

$$F(t, \tilde{x}) = \{y \in S_A^1(t, \tilde{x}) : \bar{U}(t, \tilde{x}, y) = \max_{z \in S_A^1(t, \tilde{x})} \bar{U}(t, \tilde{x}, z)\}.$$

As before, for each fixed $t \in T$, $F(t, \cdot)$ is weakly u.s.c. and for each fixed $\tilde{x} \in S_X^1$, $F(\cdot, \tilde{x})$ has a measurable graph. Furthermore, F is nonempty and convex valued. Define the correspondence $\Phi : S_X^1 \rightarrow 2^{S_X^1}$ by $\Phi(\tilde{x}) = \{\tilde{y} \in S_X^1 : \tilde{y}(t) \in F(t, \tilde{x}) \ \nu - a.e.\}$. By the u.s.c. lifting theorem (Appendix) Φ is weakly u.s.c. with convex, nonempty values (recall Aumann’s measurable selection theorem) and obviously, Φ maps points from a weakly compact (Diestel’s theorem) convex, nonempty set S_X^1 into sets of subsets of S_X^1 . By the Fan-Glicksberg fixed point theorem there exist $\tilde{x}^* \in S_X^1$ such that $\tilde{x}^* \in \Phi(\tilde{x}^*)$. It can be easily checked that \tilde{x}^* is an ex ante social equilibrium for Γ .

5.1 Convexifying effect (ex ante expected utility)

We will now show that by imposing condition (A.5), i.e., many more agents than strategies we can dispense with the concavity assumption on the utility functions and the convexity assumption on the strategy set. Thus, the counterpart of Theorem 2 for the ex ante case can be obtained.

We first need the following definition:

Definition 5.1 An *ex ante convexifying social equilibrium* for Γ is a strategy profile $\tilde{x}^* \in \int_T X$, i.e., there exist $\tilde{x}(t) \in S_{X_t}^1$, $\nu - a.e.$, $\int_T \tilde{x} = \tilde{x}^*$, such that, for $\nu - a.e.$ conditions (1) and (2) of definition 5.3. hold.

Theorem 4 Let Γ be a social system with asymmetric information satisfying (A.1), (a') and (b), (A.2), (a), (b), (d), (A.3)–(A.5). Then a convexifying ex ante social asymmetric equilibrium exists in Γ .

Proof We outline the basic argument. Define the correspondence $S_A^1 : T \times \int_T X \rightarrow 2^{L_1(\mu, Y)}$ by

$$S_A^1(t, \tilde{x}) = \{y \in S_{X_t}^1 : y(t, \omega) \in A(t, \omega, \tilde{x}) \ \mu - a.e.\}.$$

Then for each fixed $\tilde{x} \in \int_T X$, $S_A^1(\cdot, \tilde{x})$ has a measurable graph and for each fixed $t \in T$, $S_A^1(t, \cdot)$ is weakly continuous, convex. Furthermore, $S_A^1(\cdot, \cdot)$ nonempty valued. Define the set-valued function $F : T \times \int_T X \rightarrow 2^{L^1(\mu, Y)}$ by

$$F(t, \tilde{x}) = \{y \in S_A^1(t, \tilde{x}) : \bar{U}(t, \tilde{x}, y) = \max_{z \in S_A^1(t, \tilde{x})} \bar{U}(t, \tilde{x}, z)\}.$$

For each fixed $t \in T$, $F(t, \cdot)$ is weakly u.s.c., nonempty valued and for each fixed $\tilde{x} \in \int_T X$, $F(\cdot, \tilde{x})$ has a measurable graph. Define the correspondence $\Phi : \int_T X \rightarrow 2^{\int_T X}$ by $\Phi(\tilde{x}) = \int_T F(t, \tilde{x})$.

As noted earlier in the proof of Theorem 2, $\int_T X$ is weakly compact, convex, and nonempty. Furthermore, the correspondence Φ is weakly u.s.c. nonempty valued and by Theorem 7 in the Appendix, convex valued. Thus, there is a fixed point which can be easily checked that it is an ex ante convexifying social equilibrium for Γ .

6 Concluding remarks and open questions

6.1 Throughout the paper we assumed that the state space Ω is countable. The reason for this was to prove the (weak) continuity of the expected utility. It turns out that if Ω is uncountable, then the expected utility is (weakly) continuous if and only if the utility function is affine [see Balder and Yannelis (1993)]. It seems that in an expected utility set up, (either interim or ex ante) the affine linearity of the utility function is a rather strong assumption and it is less acceptable than the assumption of a countable state space.

6.2 The strategy space in this paper was assumed to be a separable Banach space. Separability seems to play an important role in the proofs as the measurable selection theorem is applied extensively. It is not clear how one can go beyond the separable case. This seems to be an open question.

6.3 Notice that the space of measure on a compact metric, endowed with the weak* topology is separable metric. Hence, if one employs weak* measurable selection theorems as well as the Gelfand integration, then the counterparts of all the theorems in this paper can be established for the space of measures. This is of interest in order to examine models of monopolistic competition or differentiated commodities.

6.4 It is well known [see for example, Arrow and Debreu (1954)], that the deterministic social equilibrium existence theorem can be used to prove the existence of a Walrasian equilibrium. One can follow the same idea to use the results of this paper to prove expected or Bayesian Walrasian equilibrium theorems for economies with asymmetric information and with a continuum of agents. It is our conjecture that, by using the theorems in this paper, the recent results of Cornet and Topuzu (forthcoming) and Balder (forthcoming) can be extended to cover asymmetric information. Moreover, the Bayesian Walrasian equilibrium existence theorems in Balder and Yannelis (submitted), Podczeck and Yannelis (submitted) can be extended to a continuum of agents. As the moment this seems to be an open question.

6.5 The dimension condition (A.5) used throughout the paper can be replaced by newer conditions which are weaker. In particular, the superatomless condition introduced by

Podczeck (2006) which is equivalent to the saturation property of a probability measure space [see Sun and Yannelis (forthcoming)], can be used to obtain Theorem 7, as well as that integration preserves u.s.c. Hence, the convexification results as well as the compactification results of this paper remain valid under the new conditions. It should be noted those conditions are not only sufficient for convexification and compactification but also necessary.

7 Appendix

The results below have been used in the proof of our main theorem. We refer the reader to Yannelis (1991) for more details and further references.

Aumann Measurable Selection Theorem *Let (T, τ, μ) be a complete finite measure space, Y be a complete, separable metric space and $\phi : T \rightarrow 2^Y$ be a nonempty valued correspondence with a measurable graph, i.e., $G_\phi \in \tau \otimes \beta(Y)$. Then there is a measurable function $f : T \rightarrow Y$ such that $f(t) \in \phi(t)$ $\mu - a.e.$*

Diestel’s Theorem *Let (T, τ, μ) be a complete finite measure space, X be a separable Banach space and $\phi : T \rightarrow 2^X$ be an integrally bounded, convex, weakly compact and nonempty valued correspondence. Then S_ϕ^1 is weakly compact in $L_1(\mu, X)$.*

Proof See Yannelis (1991).

U.S.C. Lifting Theorem *Let Y be a separable space, $(\Omega, \mathcal{F}, \mu)$ be a complete finite measure space and $X : \Omega \rightarrow 2^Y$ be an integrably bounded, nonempty, convex valued correspondence such that for all $\omega \in \Omega$, $X(\omega)$ is a compact, convex subset of Y . Denote by S_X^1 the set $\{x \in L_1(\mu, Y) : x(\omega) \in X(\omega) \mu - a.e.\}$. Let $\phi : \Omega \times S_X^1 \rightarrow 2^Y$ be a nonempty, closed, convex valued correspondence such that $\phi(\omega, x) \subset X(\omega)$ for all $(\omega, x) \in \Omega \times S_X^1$. Assume that for each fixed $x \in S_X^1$, $\phi(\cdot, x)$ has a measurable graph and that for each fixed $\omega \in \Omega$, $\phi(\omega, \cdot) : S_X^1 \rightarrow 2^Y$ is u.s.c. in the sense that the set $\{x \in S_X^1 : \phi(\omega, x) \subset V\}$ is weakly open in S_X^1 for every norm open subset V of Y . Define the correspondence $\Phi : S_X^1 \rightarrow 2^{S_X^1}$ by*

$$\Phi(x) = \left\{ y \in S_X^1 : y(\omega) \in \phi(\omega, x) \mu - a.e. \right\}.$$

Then Φ is weakly u.s.c., i.e., the set $\{x \in S_X^1 : \Phi(x) \subset V\}$ is weakly open in S_X^1 for every weakly open subset V of S_X^1 .

Proof See Yannelis (1991).

Measurability Lifting Theorem *Let Y and E be separable Banach spaces, and (T, τ, ν) and $(\Omega, \mathcal{F}, \mu)$ be finite complete measure spaces, and assume that $(\Omega, \mathcal{F}, \mu)$ is separable. Let $\gamma : T \times \Omega \times E \rightarrow 2^Y$ be a nonempty valued correspondence. Suppose that for each $y \in E$, $\gamma(\cdot, \cdot, y)$ has a measurable graph. Define the correspondence $\psi : \Omega \times E \rightarrow 2^{L_1(\mu, Y)}$ by*

$$\psi(t, y) = \{x(t) \in L_1(\mu, Y) : x(t, \omega) \in \gamma(t, \omega, y) \mu - a.e.\}.$$

Then for each $y \in E$, $\psi(\cdot, y)$ has a measurable graph.

Proof See [Balder and Yannelis \(1991\)](#).

Denote by $L_\infty(\nu)$ the space of real valued measurable, essentially bounded functions derived on (T, τ, ν) . For any $E \in \tau$ the measure space (E, τ_E, ν_E) is naturally defined, and so is the space $L_{\infty,E}(\tau) = \{f : E \rightarrow \mathcal{R}, f \text{ is } \tau_E\text{-measurable and } \nu_E\text{-essentially bounded}\}$.

For any vector space over the real field an algebraic (Hamel) basis exists. The cardinality of any Hamel basis is the same, and we denote for any vector space Y , by $\dim Y$, the cardinality of any of its bases.

Theorem A (Convexifying effect): *Let K be a weakly compact subset of a Banach space Y , and let $\varphi : T \rightarrow 2^K$ be a lower measurable closed valued correspondence. Suppose that for any pair $((T, \tau, \nu), Y)$ if $E \in \tau, \nu(E) > 0$, then $\dim L_{\infty,E}(\nu) > \dim Y$.*

Then $\int \varphi = \int \overline{\text{con}}\varphi$, (where $\overline{\text{con}}$ denotes, closed convex hull).

Proof See [\(Rustichini and Yannelis 1991, Main theorem\)](#).

The theorem above is an infinite generalization of Theorem 3 in Aumann (1964). It should be noted that the assumption that φ is lower measurable and closed valued implies that φ has a measurable graph [see for example [Castaing and Valadier \(1997\)](#)].

Define the mapping $\psi : L_1(\nu, Y) \rightarrow Y$ by $\psi(x) = \int_T x(t)d\nu(t)$, ψ is linear and norm continuous and thus weakly continuous ([Dunford and Schwartz 1958, p. 422](#)). Denote by S_φ^1 the set of all Bochner integrable selections from the correspondence $\varphi : T \rightarrow 2^Y$, i.e.,

$$S_\varphi^1 = \{y \in L_1(\nu, Y) : y(t) \in \varphi(t) \ \nu - a.e.\}.$$

The integral of φ is $\psi(S_\varphi^1) = \{\psi(x) : x \in S_\varphi^1\} = \int \varphi$. In view of the above, the reader can easily conclude that the counterparts of the u.s.c. and measurability lifting theorems for the set of all Bochner integrable selections of a correspondence, also hold for the integral of a correspondence. This is also the case for the weak compactness of the integral of a correspondence, i.e., by Diestel’s theorem⁶ it follows that $S_{\overline{\text{con}}\varphi}^1 = \{y \in L_1(\nu, Y) : y(t) \in \overline{\text{con}}\varphi(t) \ \nu - a.e.\}$ is weakly compact, and therefore, $\psi(S_{\overline{\text{con}}\varphi}^1) = \int \overline{\text{con}}\varphi$ is also weakly compact. However, by Theorem A, $\int \overline{\text{con}}\varphi = \int \varphi$, and thus $\int \varphi$ is weakly compact.

Integration preserves u.s.c. theorem: *Let (T, τ, ν) be a complete finite measure space, P be a metric space and Y be a separable Banach space. Let $\psi : T \times P \rightarrow 2^Y$ be a nonempty, compact valued correspondence such that for each fixed $t \in T$, $\psi(t, \cdot)$ is u.s.c. and that for each $p \in P$, $\psi(\cdot, p)$ has a measurable graph. Then $\int_T \psi(t, \cdot)$ is u.s.c.*

Proof See [Yannelis \(1991, Theorem 6.6 and Remark 6.1\)](#).

A version of the above theorem for weak-u.s.c. also holds, [see [\(Yannelis 1991, Theorem 5.5\)](#)].

⁶ It is assumed here that $\varphi : T \rightarrow 2^Y$ is integrably bounded and weakly compact and nonempty valued.

References

- Aliprantis, C.D., Burkinshaw, O.: Positive Operators. Academic, New York (1985)
- Arow, J., Debreu, G.: Existence of an equilibrium for a competitive economy. *Econometrica* **22**, 265–290 (1954)
- Aumann, J.R.: Integrals of set-valued functions. *J. Math. Anal. Appl.* **12**, 1–12 (1965)
- Balder, E.J.: A unifying pair of Cournot-Nash equilibrium existence results. *J. Econ. Theory* **102**, 437–470 (2002)
- Balder, E.J.: Existence of Competitive Equilibria in Economies with a Measure Space of Consumers and Consumption Externalities. *J. Math. Econ.* (forthcoming)
- Balder, E.J., Yannelis, N.C.: Equilibrium in random and Bayesian games with a continuum of players in equilibrium theory in infinite dimensional spaces. In: Khan, M.A., Yannelis, N.C. (eds.) Springer, Heidelberg (1991)
- Balder, E.J., Yannelis, N.C.: On the continuity of expected utility. *Econ. Theory* **3**, 625–643 (1993)
- Balder, E.J., Yannelis, N.C.: Bayesian Walrasian equilibrium: a new approach to rational expectations equilibrium. *Econ. Theory* (submitted)
- Castaing, C., Valadier, M.: Convex Analysis and Measurable Multifunctions, Lecture Notes in Mathematics, vol. 580. Springer, New York (1997)
- Cornet, B., Topuzu, M.: Existence of equilibrium for economies with externalities and a measure space of consumers. *Econ. Theory* (forthcoming)
- Da-Rocha, F.M., Topuzu, M.: Cournot-Nash equilibrium in continuum games with non-ordered preferences. *J. Econ. Theory* (forthcoming)
- Debreu, G.: A social equilibrium existence theorem. *Proc. Natl. Acad. Sci. USA* **38**, 803–886 (1952)
- Diestel, J., Uhl, J.: Vector measures, mathematical surveys. *Am. Math. Soc. Providence* **15**, (1977)
- Dunford, N., Schwartz, J.T.: Linear Operators, vol. I. Interscience, New York (1958)
- Haller, H.: Abstract economies with widespread externalities (1993)
- Kim, T., Yannelis, N.C.: Existence of equilibrium in Bayesian games with infinitely many players. *J. Econ. Theory* **77**, 330–353 (1977)
- Nash, J.F.: Noncooperative games. *Ann. Math.* **54**, 286–295 (1951)
- Nash, J.F.: Equilibrium points in n -person games. *Proc. Natl. Acad. Sci. USA* **36**, 48–49 (1950)
- Podczek, K.: Core and Walrasian equilibrium when agent's characteristics are extremely dispersed. *Econ. Theory* **22**, 699–725 (2003)
- Podczek, K.: On the convexity and compactness of the integral of a Banach space valued correspondence, 2006, *J. Math. Econ.* (in press)
- Podczek, K., Yannelis, N.C.: Equilibrium theory with asymmetric information and with infinitely many commodities. *J. Econ. Theory* (submitted)
- Podczek, K., Tourky, R., Yannelis, N.C.: Private expectations equilibria (2005)
- Rustichini, A., Yannelis, N.C.: What is Perfect Competition. In: Khan, M.A., Yannelis, N.C. (eds.) *Equilibrium Theory in Infinite Dimensional Spaces*, vol. 1, pp. 249–265, (1991)
- Schmeidler, D.: Equilibrium points of non-atomic games. *J. Stat. Phys.* **7**, 295–300 (1973)
- Sun Y., Yannelis, N.C.: Saturation and the integration of Banach valued correspondences, *J. Math. Econ.* (forthcoming)
- Tourky, R., Yannelis, N.C.: Markets with many more agents than commodities: Aumann's hidden assumption. *J. Econ. Theory* **101**, 189–221 (2001)
- Yannelis, N.C.: Integration of Banach-valued correspondences. In: Khan, M.A., Yannelis, N.C. (eds.) *Equilibrium Theory in Infinite Dimensional Spaces*. Springer, Heidelberg (1991)
- Yannelis, N.C.: A Bayesian equilibrium existence theorem. *Adv. Math. Econ.* **4**, 61–72 (2002)
- Yannelis, N.C., Rustichini, A.: Equilibrium points of noncooperative random and Bayesian games. In: Aliprantis, C.D. et al. (eds.) *Positive operators, Riesz spaces and economics*, vol. 2, pp. 23–48 (1991)